

## Contents of *Aquatic Botany*, Volume 74

### VOL. 74 NO. 1

SEPTEMBER 2002

Nutrient effects on autofragmentation of <i>Myriophyllum spicatum</i> D.H. Smith (Lewisville, TX, USA), J.D. Madsen (Mankato, MN, USA), K.L. Dickson and T.L. Beitingner (Denton, TX, USA) . . . . .	1
Isozyme diversity and reproduction of <i>Typha minima</i> populations on the upper River Rhine D.J. Galeuchet (Zürich, Switzerland), R. Holderegger (Birmensdorf, Switzerland), R. Rutishauser and J. Jakob Schneller (Zürich, Switzerland) . . . . .	19
Chromosomal studies on Neotropical Limncharitaceae (Alismatales) E.R. Forni-Martins and K.P. Calligaris (São Paulo, Brazil) . . . . .	33
How do water regime and clipping influence wetland plant establishment from seed banks and subsequent reproduction? K. Crosslé and M.A. Brock (Armidale, Australia) . . . . .	43
Canopy structure and pollination biology of the seagrasses <i>Posidonia australis</i> and <i>P. sinuosa</i> (Posidoneaceae) N.M. Smith and D.I. Walker (Crawley, Australia) . . . . .	57
Effects of simulated climate warming on macrophytes in freshwater microcosm communities D. McKee, K. Hatton, J.W. Eaton, D. Atkinson, A. Atherton, I. Harvey and B. Moss (Liverpool, UK) . . . . .	71
Development of amplified fragment length polymorphism markers for <i>Spartina alterniflora</i> E.J. Perkins, W.J. Streever (Vicksburg, MS, USA), E. Davis (McLean, VA, USA) and H.L. Fredrickson (Vicksburg, MS, USA) . . . . .	85

### VOL. 73 NO. 2

OCTOBER 2002

The effect of flooding regimes on decomposition of <i>Polygonum pensylvanicum</i> in playa wetlands (Southern Great Plains, USA) J.T. Anderson and L.M. Smith (Lubbock, TX, USA) . . . . .	97
Aerenchyma development and oxygen transport in the estuarine cordgrasses <i>Spartina alterniflora</i> and <i>S. anglica</i> B.R. Maricle and R.W. Lee (Pullman, WA, USA) . . . . .	109
Distribution and nutrient limitation of surfgrass, <i>Phyllospadix scouleri</i> and <i>Phyllospadix torreyi</i> , along the Pacific coast of Baja California (México) P. Ramírez-García (Coyoacán, México), J. Terrados (Blanes, Spain), F. Ramos, A. Lot, D. Ocaña (Coyoacán, México) and C.M. Duarte (Mallorca, Spain) . . . . .	121
Relationships between water level fluctuations and vegetation diversity in shallow water of New Zealand lakes T. Riis and I. Hawes (Christchurch, New Zealand) . . . . .	133

Effect of storms on photosynthesis, carbohydrate content and survival of eelgrass populations from a coastal lagoon and the adjacent open ocean A. Cabello-Pasini, C. Lara-Turrent (Baja California, Mexico) and R.C. Zimmerman (Moss Landing, CA, USA) . . . . .	149
Genetic diversity of <i>Halodule wrightii</i> using random amplified polymorphic DNA R. Angel (Houston, TX, USA) . . . . .	165

### Short communications

Lepidochronology and internodal length methods for studying <i>Posidonia oceanica</i> growth: are they compatible? A. Peirano (La Spezia, Italy) . . . . .	175
Feeding deterrence of <i>Azolla</i> in relation to deoxyanthocyanin and fatty acid composition M.F. Cohen, T. Meziane, M. Tsuchiya and H. Yamasaki (Okinawa, Japan) . . . . .	181

## VOL. 74 NO. 3

NOVEMBER 2002

Primary production and mortality of <i>Eleocharis interstincta</i> in response to water level fluctuations A. Medeiros dos Santos (Maringá, Brazil) and F. de Assis Esteves (Rio de Janeiro, Brazil) . . . . .	189
Molecular Phylogeny of Mangroves IX. Molecular marker assisted intra-specific variation and species relationships in the Indian mangrove tribe Rhizophoraceae M. Lakshmi, M. Parani and A. Parida (Chennai, India). . . . .	201
Suitability of the Fluorescence Monitoring System (FMS, Hansatech) for measurement of photosynthetic characteristics in algae J.-L. Mouget and G. Tremblin (Le Mans, France) . . . . .	219
Patterns in the secondary succession of a <i>Carex vesicaria</i> L. wetland following a permanent drawdown A. Odland (Bø, Norway) . . . . .	233
Effect of calcium, sodium and pH on uptake and accumulation of radiocesium by <i>Riccia fluitans</i> M.A. Heredia, R. Zapico, M.J. García-Sánchez and J.A. Fernández (Málaga, Spain). . . . .	245

### Short communications

A new record of a second seagrass species from the Hawaiian archipelago: <i>Halophila decipiens</i> Ostefeld K.J. McDermid, M.C. Gregoritz (Hilo, HI, USA) and D.W. Freshwater (Wilmington, NC, USA) . . . . .	257
The photosynthetic light response of <i>Halophila stipulacea</i> growing along a depth gradient in the Gulf of Aqaba, the Red Sea A.-M. Schwarz (Hamilton, New Zealand) and F. Hellblom (Stockholm, Sweden) . . . . .	263

## VOL. 74 NO. 4

DECEMBER 2002

Surface water nutrient concentrations and litter decomposition rates in wetlands impacted by agriculture and mining activities A.A. Lee and P.A. Bukaveckas (Louisville, KY, USA) . . . . .	273
Interannual and small-scale spatial variability in sexual reproduction of the seagrasses <i>Posidonia coriacea</i> and <i>Heterozostera tasmanica</i> , southwestern Australia M.L. Campey, G.A. Kendrick and D.I. Walker (Crawley, Australia) . . . . .	287
Comparison of three life-history traits of invasive <i>Elodea canadensis</i> Michx. and <i>Elodea nuttallii</i> (Planch.) H. St. John M.-H. Barrat-Segretain, A. Elger, P. Sagnes and S. Puijalon (Villeurbanne, France). . . . .	299

The effect of increasing sediment accretion on the seedlings of three common Thai mangrove species U. Thampanya (Songkhla, Thailand), J.E. Vermaat (Amsterdam, The Netherlands) and J. Terrados (Blanes, Spain) . . . . .	315
Does prolonged flooding prevent or enhance regeneration and growth of <i>Sphagnum</i> ? L. Rochefort, S. Campeau and J.-L. Bugnon (Que., Canada) . . . . .	327
Effects of ammonia in pulp mill effluents on estuarine phytoplankton assemblages: field descriptive and experimental results R.J. Livingston, A.K. Prasad, X. Niu and S.E. McGlynn (Tallahassee, FL, USA) . . . . .	343
<b>Contents of <i>Aquatic Botany</i>, Volume 74</b> . . . . .	369

the 1990s, the number of people in the world who are undernourished has increased from 600 million to 800 million.

There are a number of reasons for this increase. One of the main reasons is that the world population has increased from 5 billion in 1987 to 6 billion in 1999. This has led to a greater demand for food, which has not been met by the current level of food production.

Another reason is that the world's food production is not distributed evenly. Some countries have a surplus of food, while others have a deficit.

There are also a number of other factors that contribute to the increase in undernourishment, such as the increase in the number of people living in poverty and the increase in the number of people who are malnourished.

The World Bank has estimated that the number of people who are undernourished will increase to 1 billion by the year 2020.

This is a very worrying trend, and it is important that we take action to address this problem.

There are a number of ways in which we can address this problem. One of the most important is to increase the world's food production.

This can be done by increasing the area of land that is used for agriculture, by increasing the productivity of the land that is already being used, and by increasing the number of people who are involved in food production.

Another way to address the problem is to improve the distribution of food. This can be done by increasing the number of people who are involved in food distribution, and by improving the infrastructure that is used for food distribution.

There are also a number of other ways in which we can address the problem, such as by increasing the number of people who are involved in food production and distribution, and by improving the infrastructure that is used for food production and distribution.

It is important that we take action to address this problem, as it is a major threat to the world's food security.

The World Bank has estimated that the number of people who are undernourished will increase to 1 billion by the year 2020.

This is a very worrying trend, and it is important that we take action to address this problem.

There are a number of ways in which we can address this problem. One of the most important is to increase the world's food production.

This can be done by increasing the area of land that is used for agriculture, by increasing the productivity of the land that is already being used, and by increasing the number of people who are involved in food production.

Another way to address the problem is to improve the distribution of food. This can be done by increasing the number of people who are involved in food distribution, and by improving the infrastructure that is used for food distribution.

There are also a number of other ways in which we can address the problem, such as by increasing the number of people who are involved in food production and distribution, and by improving the infrastructure that is used for food production and distribution.

It is important that we take action to address this problem, as it is a major threat to the world's food security.

The World Bank has estimated that the number of people who are undernourished will increase to 1 billion by the year 2020.

This is a very worrying trend, and it is important that we take action to address this problem.

There are a number of ways in which we can address this problem. One of the most important is to increase the world's food production.

This can be done by increasing the area of land that is used for agriculture, by increasing the productivity of the land that is already being used, and by increasing the number of people who are involved in food production.

Another way to address the problem is to improve the distribution of food. This can be done by increasing the number of people who are involved in food distribution, and by improving the infrastructure that is used for food distribution.

There are also a number of other ways in which we can address the problem, such as by increasing the number of people who are involved in food production and distribution, and by improving the infrastructure that is used for food production and distribution.

It is important that we take action to address this problem, as it is a major threat to the world's food security.

The World Bank has estimated that the number of people who are undernourished will increase to 1 billion by the year 2020.

This is a very worrying trend, and it is important that we take action to address this problem.

